

Modal Propositional Logic

David Pearce

Madrid, November 2006

Modal propositional logic: Syntax

- Syntax of classical propositional logic
- Countable set of propositional atoms: $p_0, p_1, \dots, p_n, \dots$
- Logical connectives: $\wedge, \vee, \rightarrow, \neg$
- Modal operators: \Box, \Diamond (necessity, possibility)
- If α is a wff, so is $\Box\alpha$. $\Diamond\alpha =_{df} \neg\Box\neg\alpha$

Normal Modal Systems

- A *system of modal logic* is a certain class S of formulas whose elements are *theorems*
- $\vdash_S \alpha$ denotes that α is a theorem of S .
- A modal system is *normal* if it contains:
 - all theorems of propositional logic
 - the axiom **K**: $\Box(p \rightarrow q) \rightarrow (\Box p \rightarrow \Box q)$
 - the rules US (uniform substitution): if α is a theorem so is any substitution instance
 - MP (modus ponens)
 - N (necessitation): $\vdash \alpha \Rightarrow \vdash \Box \alpha$

Models of normal systems

A model is a triple $\langle W, R, V \rangle$, where

- W is a non-empty set (of possible worlds)
- R is a binary relation over W , ie $R \subseteq W \times W$
- V is a valuation assigning a truth-value 1 or 0 to each atomic proposition p at each world $w \in W$

Models of normal systems, contd

Valuations V are extended to all formulas via the following rules:

$$(\mathbf{V} \wedge) \quad V(\alpha \wedge \beta, w) = 1 \text{ iff } V(\alpha, w) = 1 \ \& \ V(\beta, w) = 1$$

$$(\mathbf{V} \vee) \quad V(\alpha \vee \beta, w) = 1 \text{ iff } V(\alpha, w) = 1 \text{ or } V(\beta, w) = 1$$

$$(\mathbf{V} \rightarrow) \quad V(\alpha \rightarrow \beta, w) = 1 \text{ iff } V(\alpha, w) = 1 \text{ implies } V(\beta, w) = 1$$

$$(\mathbf{V} \neg) \quad V(\neg\alpha, w) = 1 \text{ iff } V(\alpha, w) = 0$$

$$(\mathbf{V} \Box) \quad V(\Box\alpha, w) = 1 \text{ iff } V(\alpha, w') = 1, \text{ for all } w' \text{ such that } wRw'$$

Truth and Validity

- A formula α is true at world w in a model $\mathcal{M} = \langle W, R, V \rangle$ if $V(\alpha, w) = 1$. In this case we sometimes write $\mathcal{M}, w \models \alpha$.
- A formula α is true in a model $\langle W, R, V \rangle$ if $V(\alpha, w) = 1$, for all $w \in W$. We also write $\mathcal{M} \models \alpha$.
- A formula is valid (in a class of models) if it is true in every model in that class
- The axiom **K** is valid in the class of all models
- The weakest normal system axiomatised by **K** and propositional logic is called K . Its theorems are true in all models

some other axioms of normal systems

Stronger normal systems can be obtained by adding further axioms

$$\mathbf{T} : \Box p \rightarrow p$$

$$\mathbf{D} : \Box p \rightarrow \Diamond p$$

$$\mathbf{4} : \Box p \rightarrow \Box \Box p$$

$$\mathbf{5} : \Diamond \Box p \rightarrow \Box p$$

$$\mathbf{B} : \Diamond \Box p \rightarrow p$$

$$\mathbf{W5} : \Diamond \Box p \rightarrow (p \rightarrow \Box p)$$

$$\mathbf{F} : (p \wedge \Diamond \Box q) \rightarrow \Box (\Diamond p \vee q)$$

some normal systems

Some well-known normal systems are denoted as follows

$B : \mathbf{B}$

$T : \mathbf{K, T}$

$S4 : \mathbf{K, T, 4}$

$S4F : \mathbf{K, T, 4, F}$

$KD45 : \mathbf{K, D, 4, 5}$

$SW5 : \mathbf{K, T, 4, W5}$

$S5 : \mathbf{K, T, 4, 5}$

Some relations between normal systems

□ $K \subset T \subset S4 \subset S5$

□ $K \subset B \subset S4 \subset S5$

□ These logics are *sound* with respect to classes of models whose accessibility relations satisfy simple algebraic properties.

Some types of binary relations

Let R be a binary relation over a set X .

- R is *reflexive* if $R(a, a)$ for every $a \in X$
- R is *symmetric* if $R(a, b) \Rightarrow R(b, a)$ for every $a, b \in X$
- R is *transitive* if $R(a, b), R(b, c) \Rightarrow R(a, c)$ for every $a, b, c \in X$
- A relation that is reflexive, symmetric and transitive is said to be an *equivalence* relation.

Some soundness characterisations

- T is sound for the class of reflexive models, ie. the axiom $\mathbf{T} : \Box P \rightarrow p$ is valid in models whose R -relation is reflexive.
- $S4$ is sound wrt to models that are reflexive and transitive
- B is sound wrt models that are reflexive and symmetric.
- $S5$ is sound wrt models in which R is an equivalence relation.

soundness proofs

To prove soundness we must show

- The axioms of the system are true in all models of the given class
- The transformation rules US, MP and N are truth preserving, ie when applied to formulas true in all models, they lead to formulas true in all models.

soundness for K

so to prove soundness for the system K we must show

- The **K** axiom is true in all models. Given a model $\langle W, R, V \rangle$, it suffices to show that if (a) $\Box(p \rightarrow q)$ and $\Box p$ are true in a world w , then also (b) $\Box q$ is true in w . Suppose (a) holds. Then by $(V \Box)$, $p \rightarrow q$ and p are true in all w' such that $R(w, w')$, so by $(V \rightarrow)$ so is q . Therefore by $(V \Box)$, $\Box q$ is true in w .
- The transformation rules US, MP and N are validity preserving, ie when applied to formulas true in all models, they lead to formulas true in all models. Suppose α is valid, then it is a formula true in every world w in any model. Then α is true independent of the truth-values assigned to the atomic variables in α . Hence if β is the result of uniformly replacing the variables of α by any wff, then β must also be true in w . So the rule US is validity preserving.

Exercises

- (1) show that the rules **MP** and **N** are validity preserving.
- (2) show that the axiom **T** is true in all reflexive models.
- (3) show that the axiom **B** is true in all reflexive, symmetric models.

preserving validity in a (single) model

The rule US of uniform substitution does not preserve truth in a single model. Counter-example: consider a model with two worlds w, w' with (w, w') as the only element in the R relation. Consider atoms p, q where p is true at both worlds and q at just the world w' . Then $\Box p \rightarrow p$ is true in the model but $\Box q \rightarrow q$ is not. Yet the latter is a substitution instance of the former.

However we do have the following:

- Theorem. Let S be an axiomatic, normal model system and let $\langle W, R, V \rangle$ be any model. If every substitution instance of every axiom of S is true in $\langle W, R, V \rangle$, then every theorem of S is true in $\langle W, R, V \rangle$.
- Note that the rules MP and N do preserve truth in a single model.

more on binary relations

Let R be a binary relation over a set X .

- R is *universal* if $R = X \times X$
- R is *Euclidean* if for every $a, b, c \in X$ such that $R(a, b)$ and $R(a, c)$, also $R(b, c)$
- Suppose $a \in X$ and there is no $b \in X$ such that $R(a, b)$, then a is called a *dead-end*.
- Note that if w is a dead-end, then we always have $V(\Box\alpha, w) = 1$ and $V(\Diamond\alpha, w) = 0$

completeness via canonical models

We want to show that certain classes of models *fully* characterise particular normal model systems. We use the powerful method of *canonical* models.

- Let S be a normal modal system and C a given class of models. A wff is said to be C -valid iff it is true in every model in C .
- S is *sound* wrt C if every theorem of S is C -valid.
- S is *complete* wrt C if every C -valid formula is a theorem of S ; ie. if α is not a theorem of S ($\not\vdash_S \alpha$) then it is not true in some C -model.
- A formula α is said to be *S -inconsistent* if $\vdash_S \neg\alpha$; otherwise (if $\not\vdash_S \neg\alpha$) it is *S -consistent*. It follows that S is complete wrt C if $\forall\alpha$, if α is S -consistent then there is a C -model in which α is true at some world w .

maximal consistent sets

We first generalise S -consistency to sets of formulas

- Definition: A finite set $\Sigma = \{\alpha_1, \dots, \alpha_n\}$ is S -consistent iff $\alpha_1 \wedge \dots \wedge \alpha_n$ is S -consistent.
- An arbitrary set of formulas Σ is S -consistent if every finite subset of Σ is S -consistent, ie there is no finite $\{\alpha_1, \dots, \alpha_n\} \subset \Sigma$ such that $\vdash_S \neg(\alpha_1 \wedge \dots \wedge \alpha_n)$.
- the canonical model method will show that if Σ is an S -consistent set of wff, then there is a C -model $\mathcal{M} = \langle W, R, V \rangle$ and $w \in W$ such that $\mathcal{M}, w \models \Sigma$. \mathcal{M} is called the canonical model.

maximal consistent sets, contd

□ Definition: A set Γ of wff is maximal iff for every wff α , either $\alpha \in \Gamma$ or $\neg\alpha \in \Gamma$.

□ Γ is said to be *maximal S-consistent* iff it is maximal and *S-consistent*.

Lemma 1 Let Γ be a maximal *S-consistent* set of wff. Then:

1. for any α , exactly one member of $\{\alpha, \neg\alpha\}$ is in Γ
2. $\alpha \vee \beta \in \Gamma$ iff either $\alpha \in \Gamma$ or $\beta \in \Gamma$
3. $\alpha, \beta \in \Gamma$ iff both $\alpha \in \Gamma$ and $\beta \in \Gamma$
4. $\vdash_S \alpha \Rightarrow \alpha \in \Gamma$
5. if $\alpha \in \Gamma$ and $\alpha \rightarrow \beta \in \Gamma$, then $\beta \in \Gamma$
6. if $\alpha \in \Gamma$ and $\vdash_S \alpha \rightarrow \beta$, then $\beta \in \Gamma$

□ **Theorem 2** Let Σ be *S-consistent*. Then there is a maximal *S-consistent* set $\Gamma \supseteq \Sigma$.

canonical models

First some notation: for Σ a set of wff, let $\Box^-(\Sigma) =_{df} \{\alpha : \Box\alpha \in \Sigma\}$

- **Lemma 3** let S be a normal system and Γ an S -consistent set of wff containing a wff of the form $\neg\Box\alpha$. Then $\Box^-(\Sigma) \cup \{\neg\alpha\}$ is S -consistent.
- **Corollary.** let S be normal and Γ an S -consistent set of wff containing a wff of the form $\Diamond\alpha$. Then $\Box^-(\Sigma) \cup \{\alpha\}$ is S -consistent.
- **Definition.** The *canonical* model of a normal modal system S is the model $\langle W, R, V \rangle$ defined as follows.
 1. $W = \{w : w \text{ is a maximal } S\text{-consistent set of wff}\}$
 2. For any $w, w' \in W$, $R(w, w') \Leftrightarrow \Box^-(w) \subseteq w'$.
 3. for any atom p and $w \in W$, $V(p, w) = 1 \Leftrightarrow p \in w$.

basic theorem for canonical models

□ **Theorem 4** Let $\langle W, R, V \rangle$ be the *canonical* model of a normal modal system S . For any wff α and any $w \in W$, $V(\alpha, w) = 1 \Leftrightarrow \alpha \in w$.

□ **Proof.** By induction on complexity of α

1. For α an atom, claim holds by definition.
2. Assume theorem for α and prove for $\neg\alpha$. Consider any $\neg\alpha$ and $w \in W$. By $(V \neg)$, $V(\neg\alpha, w) = 1 \Leftrightarrow V(\alpha, w) = 0$. By assumption, $V(\alpha, w) = 0 \Leftrightarrow \alpha \notin w$. Hence $V(\neg\alpha, w) = 1 \Leftrightarrow \alpha \notin w$. By Lemma 1.1, $\alpha \notin w$ iff $\neg\alpha \in w$. Therefore $V(\neg\alpha, w) = 1 \Leftrightarrow \neg\alpha \in w$.
3. For $\alpha \vee \beta$, assume claim holds for α and β , use $(V \vee)$ and apply Lemma 1.2.
4. Consider the case of $\Box\alpha$ and assume claim holds for α . (i) suppose $\Box\alpha \in w$. By definition of R , $\alpha \in w'$ for all w' such that $R(w, w')$. By induction assumption, for each such w' , $V(\alpha, w') = 1$. So by $(V \Box)$, $V(\Box\alpha) = 1$.

(ii) Suppose on the other hand that $\Box\alpha \notin w$. By Lemma 1.1, $\neg\Box\alpha \in w$. So by Lemma 3, $\Box^-(w) \cup \{\neg\alpha\}$ is S -consistent. Thus by Theorem 2 and definition of W , there exists a $w' \in W$ such that $\Box^-(w) \cup \{\neg\alpha\} \subseteq w'$. Hence we have (i) $\Box^-(w) \subseteq w'$ and (ii) $\neg\alpha \in w'$. (i) implies $R(w, w')$, by def of R . So by induction assumption, theorem holds for α and by part 1 above for $\neg\alpha$. Therefore by (ii), since $\neg\alpha \in w'$, we have $V(\neg\alpha, w') = 1$ and therefore $V(\alpha, w') \neq 1$. Then by $(V \Box)$, we obtain $V(\Box\alpha) \neq 1$.

□ Corollary. A formula α is valid in the canonical model for S iff $\vdash_S \alpha$. Proof: Let $\langle W, R, V \rangle$ be the *canonical* model for S . Suppose that $\vdash_S \alpha$. Then by Lemma 1.4, α belongs to every maximal S -consistent set. So $\alpha \in w$, for all $w \in W$. By Theorem 4, $V(\alpha, w) = 1$, for all $w \in W$ so α is true in the canonical model. Suppose that $\not\vdash_S \alpha$. Then $\neg\alpha$ is S -consistent. So for some $w \in W$, $\neg\alpha \in w$ and hence $\alpha \notin w$. Therefore by Theorem 4, $V(\alpha, w) \neq 1$, for some $w \in W$, and so α is not true in the canonical model.

□ Corollary. The system K is complete for the class of all models.

some completeness theorems

- T is complete with respect to the class of all reflexive models
- $S4$ is complete for the class of all reflexive, transitive models
- B is complete for the class of all reflexive, symmetrical models
- $S5$ is complete for the class of all models in which R is an equivalence relation

Method Show in each case that the canonical model has the stated structure.

more completeness results

- D is complete with respect to the class of all models with serial accessibility relation
- $KD45$ is complete for the class of all transitive, Euclidean models with no dead-ends
- $S4F$ is complete for the class of all reflexive, transitive models with the condition: if $R(a, b)$ and $R(a, c)$ but not $R(b, a)$, then $R(c, b)$.
- $SW5$ is complete for the class of all reflexive, transitive models in which R satisfies the condition: if $a \neq b, a \neq c, R(a, b)$ and $R(a, c)$, then $R(b, c)$ and $R(c, b)$.

References

G E Hughes & M J Cresswell, *A Companion to Modal Logic*, Methuen, 1984.

V W Marek & M Truszczyński, *Nonmonotonic Logic*, Springer, 1993

W. Rautenberg, *Klassische und nichtklassische Aussagenlogik*, Vieweg, 1979